

Evidence-Based Care Measures 2011

Breast Cancer Screening	Women 50-69 years old who have had a mammogram during the measurement period or the year prior
Cervical Cancer Screening	Women 24-64 years old who had a Pap during the measurement year or the year prior
Appropriate Treatment of Children with URI	Children 3 months to 18 years old who were given a diagnosis of URI and were not dispensed an antibiotic prescription
Antibiotics: Adults with Acute Bronchitis	Adults 18-64 with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription
Asthma	Patients 5-50 years old who were identified as having persistent asthma and were appropriately prescribed medication during the measurement period
Adolescent Well Care Visits	Patients 12-21 years old who have had at least one visit with a PCP or OB/GYN practitioner during the measurement period
Adolescent Immunization	Patients turning 13 during the measurement period who have had one dose of meningococcal vaccine and one Tdap or one Td by their birthday
Childhood Immunization Status	Patients who have received four DTaP, three IPV, one MMR, two HiB, three HepB, one VZV, four PCV, two HepA, two or three RV and two flu vaccines by their second birthday
Well Child Visits	Patients who turned 15 months old during the measurement period who have had 5 or more well child visits
Older Well Child Visits	Patients 3-6 years old who have had at least one well child visit during the measurement period
Spirometry and COPD	Patients with a new diagnosis of COPD who received a spirometry test to confirm diagnosis
CHF LDL Screening	Patients 18-75 years of age with congestive heart failure who had an LDL-C test
CHF ACE/ARB	Patients with a diagnosis of CHF who received ACE/ARB therapy during the measurement period
Beta-Blocker After Acute MI	Patients 18 and older who were hospitalized with an acute MI who received beta-blocker treatment for 6 months after discharge
CAD LDL-C	Patients 18-75 with CAD who had an LDL-C screening

Evidence-Based Care Measures 2011

CAD-Lipid Lowering Drugs	Patients between 18 and 75 years old with Coronary artery disease who received a lipid lowering drug
CAD-Statin Use	Patients 18-75 with a diagnosis of CAD who have at least one statin prescription filled during the measurement year
AF and Flutter-anticoagulation	Patients over 18 with a diagnosis of non-valvular AF or flutter at high risk for thromboembolism who were prescribed warfarin
AF and Flutter INR	Patients 18-64 with AF and Flutter on warfarin who received INR at least monthly
CAD-Cardiac Rehabilitation	Patients 18-64 that have been discharged from hospitalization who have had a qualifying event/dx who have attended a cardiac rehab program
Diabetes HbA1c	Patients 18-75 with diabetes who had an HbA1c test
Diabetes LDL-C	Patients 18-75 with diabetes who had a LDL-C screening
Diabetes monitor nephropathy	Patients 18-75 with diabetes who had a nephropathy screening test
Diabetes lipid lowering drug	Patients 40-75 with diabetes who received a prescription for at least one lipid lowering drug
Diabetes-Statin	Patients 40-75 with diabetes who have at least one statin prescription
Diabetes/CHF-ACE/ARB	Patients 18-75 with diabetes and comorbid CHF who received ACE/ARB therapy
Diabetes/Nephropathy ACE/ARB	Patients 18-75 with diabetes and comorbid nephropathy who received ACE/ARB therapy
Diabetes/HPTN ACE/ARB	Patients 18-75 with diabetes and comorbid hypertension who received ACE/ARB therapy
Low Back Pain	Patients 18-50 with a diagnosis of low back pain who did not have an imaging study in the 28 days following the initial diagnosis
Antidepressant Med	Patients 18 and older who were diagnosed with a new episode of major depression remained on antidepressant medication at least 12 weeks for the acute period and 6 months for the continuation phase
Medication Management	Annual monitoring for patients 18 and older on persistent medications