



One of a series of tip sheets that look at key Healthcare Effectiveness Data and Information Set measures

Flu vaccinations for adults ages 18-64

This measure, also referred to as FVA, examines the percentage of patients ages 18 to 64 who received an influenza vaccination during the prior flu season.

Exclusions:

Exclude all patients with an allergy to eggs or history of a serious adverse or allergic reaction to an influenza vaccine at any time prior to or during the measurement year.

Improving HEDIS® scores*

Proper documentation is critical:

- Provide clear documentation of vaccines you administered or documentation of vaccines received elsewhere, with dates and locations.
- Utilize the Michigan Care Improvement Registry, or MCIR, tracking system for all patients.
- Strongly recommend that patients get a flu vaccine and educate them about vaccine efficacy and safety.
- Influenza often leads to hospitalization for patients with chronic or underlying conditions. Consider flagging charts for high-risk patients and sending flu vaccine reminders by mail, email or text.
- Keep in mind that flu shot compliance is obtained through the annual Consumer Assessment of Healthcare Providers and Systems, or CAPHS, survey. It depends on patient recollection of receiving the vaccination.

Did you know?

- Each year in the U.S., 5 to 20 percent of the population is infected with the flu.
- More than 200,000 people are hospitalized annually in the U.S. for complications from the flu.
- From 1980-2010, flu-associated deaths in the U.S. ranged from 3,000 to 49,000 annually.

Tip:

Make vaccine status assessment and discussion an integral part of every office visit.

*HEDIS, which stands for Healthcare Effectiveness Data and Information Set, is a registered trademark of the National Committee for Quality Assurance.